

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO

Molecular identification of *Mycobacterium tuberculosis* complex by region of differentiation-typing and polymerase chain reaction-restriction fragment length polymorphism method

Seydeh Zeinab Mirzaki^a, Nader Mosavari^{b,*}, Razieh Nazari^a, Morteza Akbarian^b

^a Department of Microbiology, Qom Branch, Islamic Azad University, Qom, Iran

^b Bovine Tuberculosis Reference Laboratory, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran

ARTICLE INFO

Article history:

Received 25 September 2016

Accepted 1 October 2016

Available online xxxx

Keywords:

IS6110

Mycobacterium tuberculosis complex

PCR-16srRNA

PCR-RFLP

RD-typing

ABSTRACT

Tuberculosis (TB) is one of the most common zoonotic infectious diseases in the world. Identification of *Mycobacterium* isolates is essential for proper treatment of TB. The aim of this study was to identify *Mycobacterium* isolates collected from TB patients in Alborz Province, Iran, by region of differentiation (RD)-typing. Fifty samples from tuberculosis patients were cultured in pyruvate and glycerinated Lowenstein-Jensen medium. DNA was extracted from the isolates by the van Solingen method and subjected to polymerase chain reaction (PCR)-16SrRNA, PCR-IS6110, and RD-typing with primers RD1, RD4, RD9, and RD12, respectively. Out of 50 isolates, only one isolate appeared negative in IS6110-PCR and was considered nontuberculosis complex. The remaining isolates gave PCR products of approximately 543 bp, 245 bp, 146 bp, 172 bp, 235 bp, and 369 bp with 16s-rRNA, IS6110-PCR, RD-1, RD-4, RD-9, and RD-12 PCR, respectively. PCR-restriction fragment length polymorphism of *oxyR* pseudogene confirmed the results. All isolates except one from Alborz Province appeared positive for *Mycobacterium tuberculosis*. Based on the obtained results, all isolates except one were identified as *M. tuberculosis*. The only negative isolate appeared 93% and 97% similar to *Nocardia* or *Mycobacterium* sp. (*Mycobacterium neoaurum*), respectively, based on sequencing and alignment of 16s-rRNA and *hsp65*. Accurate identification of *Mycobacterium* isolates is of utmost importance for proper and immediate treatment of TB patients. In this study, RD-typing appeared to be a suitable method for correct identification of *M. tuberculosis* isolates.

Conflicts of interest

The authors declare no conflicts of interest.

* Corresponding author at: Bovine Tuberculosis Reference Laboratory, Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Tehran, Iran.

E-mail address: nmosavari@gmail.com (N. Mosavari).

Peer review under responsibility of Asian African Society for Mycobacteriology.

<http://dx.doi.org/10.1016/j.ijmyco.2016.10.004>